Measurement and simulation of volatile particle emissions from military aircraft

WP 1625, WP-1626, WP-1627

Allen Robinson
Carnegie Mellon University

Presented at Partners in Environmental Technology Technical Symposium & Workshop Washington DC November 30, 2011.



Report Documentation Page

Form Approved OMB No. 0704-0188

Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.

1. REPORT DATE NOV 2011	2. REPORT TYPE	3. DATES COVERED 00-00-2011 to 00-00-2011	
4. TITLE AND SUBTITLE	5a. CONTRACT NUMBER		
Measurement and simulation of volatile particle emissions from military aircraft		5b. GRANT NUMBER	
		5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)	5d. PROJECT NUMBER		
	5e. TASK NUMBER		
	5f. WORK UNIT NUMBER		
7. PERFORMING ORGANIZATION NAME(S) AND ALL Carnegie Mellon University,5000 Forb	8. PERFORMING ORGANIZATION REPORT NUMBER		
9. SPONSORING/MONITORING AGENCY NAME(S) A	10. SPONSOR/MONITOR'S ACRONYM(S)		
	11. SPONSOR/MONITOR'S REPORT NUMBER(S)		

12. DISTRIBUTION/AVAILABILITY STATEMENT

Approved for public release; distribution unlimited

13. SUPPLEMENTARY NOTES

Presented at the Partners in Environmental Technology Technical Symposium & Workshop, 29 Nov? 1 Dec 2011, Washington, DC. Sponsored by SERDP and ESTCP. U.S. Government or Federal Rights License

14. ABSTRACT

Many military airfields are located in nonattainment areas for fine particulate matter (PM2.5) but the contribution of aircraft emissions to local and regional PM2.5 concentrations is not well understood. Aircraft directly emit particles, but the vast majority of the emissions are gases and vapors, some of which undergo gas-to-particle conversion in the atmosphere. Primary particulate matter is defined as directly emitted particles plus any material that condenses into the particle phase without undergoing chemical reactions. This latter component is often referred to as volatile particulate matter. Secondary particulate matter is formed from oxidation of gas-phase species such as sulfur dioxide and organics. To develop effective control strategies one must understand the overall contribution of military aircraft emissions to ambient particulate matter? both direct particle emissions and particulate matter formed in the atmosphere. To assist the Department of Defense in meeting the current and future regulations related to PM2.5 concentrations, SERDP initiated a research program in 2007 to investigate volatile particulate matter emissions. Three projects were funded within this program; the lead organizations of these programs were Aerodyne Research Inc., Carnegie Mellon University, and Oak Ridge National Laboratories. The projects included experiments to measure the formation and atmospheric evolution of volatile particulate matter, the development volatile emission models, and the evaluation of new instrumentation and experimental techniques. This presentation will summarize the major findings of the SERDP volatile particulate matter program, including the contribution of lubricating oil and atmospheric oxidation to volatile particulate matter emissions.

15. SUBJECT TERMS

16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF	18. NUMBER	19a. NAME OF
			ABSTRACT	OF PAGES	RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified	Same as Report (SAR)	26	1.00.01.01.02.01.01.00.01

Standard Form 298 (Rev. 8-98) Prescribed by ANSI Std Z39-18

MEASUREMENT AND SIMULATION OF VOLATILE PARTICLE EMISSIONS FROM MILITARY AIRCRAFT

DR. ALLEN ROBINSON Carnegie Mellon University 5000 Forbes Avenue Pittsburgh, PA 15213 (412) 368-3657 alr@andrew.cmu.edu


Many military airfields are located in nonattainment areas for fine particulate matter (PM_{2.5}), but the contribution of aircraft emissions to local and regional PM_{2.5} concentrations is not well understood. Aircraft directly emit particles, but the vast majority of the emissions are gases and vapors, some of which undergo gas-to-particle conversion in the atmosphere. Primary particulate matter is defined as directly emitted particles plus any material that condenses into the particle phase without undergoing chemical reactions. This latter component is often referred to as volatile particulate matter. Secondary particulate matter is formed from oxidation of gas-phase species such as sulfur dioxide and organics. To develop effective control strategies one must understand the overall contribution of military aircraft emissions to ambient particulate matter—both direct particle emissions and particulate matter formed in the atmosphere.


To assist the Department of Defense in meeting the current and future regulations related to $PM_{2.5}$ concentrations, SERDP initiated a research program in 2007 to investigate volatile particulate matter emissions. Three projects were funded within this program; the lead organizations of these programs were Aerodyne Research Inc., Carnegie Mellon University, and Oak Ridge National Laboratories. The projects included experiments to measure the formation and atmospheric evolution of volatile particulate matter, the development volatile emission models, and the evaluation of new instrumentation and experimental techniques. This presentation will summarize the major findings of the SERDP volatile particulate matter program, including the contribution of lubricating oil and atmospheric oxidation to volatile particulate matter emissions.



What is volatile PM?










How does volatile PM evolve in the atmosphere?



Spatial / Temporal Scales




SERDP Volatile PM Projects

Oak Ridge National Lab Team (WP-1627)

- Measurement Systems
 - Thermodenuder
 - Dilution sampler & aging chamber
- Measurements of gas turbine engine emissions







Aerodyne Team (WP-1625)

- Microphysical Kinetic Model for Plume
- Contributions of lube oil
- Laboratory studies of specific HCs and Sulfate

 $0 \rightarrow 1 \text{ km}$

Combustor system studies










$0.1 \rightarrow 50 \text{ km}$

Carnegie Mellon University Team (WP-1626)

- Measurement of PM in engine exhaust
- Aging in "smog" chamber
- Modeling of gas-particle partitioning and SOA production




Microphysical Modeling of Near-Field Evolution of Volatile PM






Predicted evolution of nucleation mode




Distance downwind (km)



Predicted Particle Composition in Aircraft Plume







Findings:


- Highly Dynamic Systems
- Nucleation driven by sulfuric acid and water
- Hydrocarbons critical for growth





Staged Testing





















What are effects of photochemical aging?










Taking off the chamber cover

<u>WP-1626</u>




Primary PM in Chamber



17




Secondary aerosol production overwhelms primary emissions




SERDP DOE

At every engine load





Effect of Fuel Composition on Secondary Aerosol






Modeling Secondary Aerosol Formation

Parameterizing Yields

Predicted Evolution Yields




Secondary PM dominant at urban and regional scales


WP-1626₁



Instrumentation development

Dilution sampler






New Thermodenuder



- Low particle losses
- Portability
- Turnkey



Lubrication Oil and Volatile PM



Findings:

The emission of lubrication oil varies significantly among the aircraft engines with different designs

Contribution from oil can be dominant in organic PM emission in some cases.




Key findings SERDP Volatile PM

- Volatile PM highly dynamic system in atmosphere
- Secondary volatile PM production very significant
- Lube oil important constituent of volatile PM
- New tools:
 - Detailed microphysical kinetic model for plume soot, sulfate, and organic
 - Volatility basis set model for gas-particle partitioning of primary organic aerosol and production of secondary organic aerosol (SOA) in atmosphere
 - Dilution sampler, thermodenuder, and smog chamber techniques
- Archival Papers: 7 published, 4 near submission, others in process



Primary PM Emissions

CFM56 with Engine Load



T63 with Fuel Composition

